



Stop a stroke in its tracks



Strokes cause damage to the brain by limiting or stopping blood flow, preventing oxygen and nutrients from getting to brain cells.¹ Any stroke is serious, and can cause brain damage or death. But getting help quickly can give you a better shot at making a full recovery.²

Prevention

Your risk of having a stroke goes up as you age. The good news is, there are things you can do to lower your risk.³

- Don't smoke.
- Keep your blood pressure and cholesterol levels healthy.
- Lose weight, if needed.
- Eat a well-balanced, nutritious diet.
- Exercise regularly.
- Limit alcohol to one (for women) or two (for men) drinks a day.
- Take preventive medication like aspirin, if your doctor suggests it.

Know the signs of stroke

Getting help quickly after a stroke can be the key to preventing death or serious disability. The best treatments only work if they are given within three hours of the first symptom. That's why it's so important to know the signs that someone is having a stroke.

- Sudden **weakness or numbness** in the face, arm or leg, especially on only one side of the body.
- Sudden **confusion** or trouble talking.
- Sudden **trouble walking**, dizziness or loss of balance.
- Sudden **severe headache**.
- Sudden **trouble seeing** out of one or both eyes.

If you're not sure whether someone is having a stroke, think F.A.S.T.²

F – Face: Ask the person to smile. Does one side of their face droop?

A – Arms: Ask them to raise both arms. Does one arm drift downward?

S – Speech: Ask them to repeat a simple phrase. Does their speech sound slurred?

T – Time: If you see any of the signs, even if they start to go away, call 9-1-1 for an ambulance. Paramedics can start life-saving treatment on the way to the emergency room.

This information is meant to be educational. It should not be interpreted as medical advice. Please talk to your doctor about changes that may affect your health.



Sources:

1 Centers for Disease Control and Prevention website: Stroke Facts (accessed February 2017); www.cdc.gov/stroke/facts.htm.

2 Centers for Disease Control and Prevention website: Stroke Signs and Symptoms (accessed February 2017); www.cdc.gov/stroke/signs_symptoms.htm.

3 Centers for Disease Control and Prevention website: Preventing Stroke: Healthy Living (accessed February 2017); www.cdc.gov/stroke/healthy_living.htm.

4 American Stroke Association website: Understanding Risk (accessed February 2017); www.strokeassociation.org.

5 Centers for Disease Control and Prevention website: Sickle Cell Disease (SCD) (accessed February 2017); www.cdc.gov/ncbddd/sicklecell/data.html.

6 Centers for Disease Control and Prevention website: Recovering from Stroke (accessed February 2017); www.cdc.gov/stroke/recovery.htm.

Anthem Blue Cross is the trade name of Blue Cross of California. Anthem Blue Cross and Anthem Blue Cross Life and Health Insurance Company are independent licensees of the Blue Cross Association. ANTHEM is a registered trademark of Anthem Insurance Companies, Inc.

35986CAMENABC VPOD Rev. 05/17



Know your risk factors

A risk factor is anything that raises your chance of having a stroke. Making healthy choices like managing your weight, eating right, staying active, and not smoking can help lower your risk. Other risk factors are out of your control:^{4,5}

- **Age** – You can have a stroke at any age, but your risk goes up as you get older.
- **Gender** – Men are more likely to have a stroke than women.
- **Health history** – If you or your family members have sickle cell disease, diabetes, high blood pressure, high cholesterol or heart problems, that raises your risk. Being overweight or having had a previous stroke or Transient Ischemic Attack (TIA) can also make you more likely to have a stroke in the future. Work closely with your doctor to manage your health conditions.

Treatment

People who arrive at the hospital within three hours of their first symptom are more likely to make a full recovery.¹ That's why it's so important to get emergency care at the first sign of a stroke.

A stroke can be treated with drugs or surgery. As a person recovers, they may need:⁶

- Physical therapy, to help with movement and balance.
- Occupational therapy, to relearn things like eating, drinking, reading or getting dressed.
- Speech therapy.
- Mental health therapy, to help with depression and trouble with attention, expressing emotions and other conditions that are common after a stroke.

